



ALBORVET

A BROAD SPECTRUM ANTHELMINTIC FOR LIVESTOCK AND POULTRY

MICRONIZED

COMPOSITION

Each bolus contains:
Albendazole IP
4000 mg, 3000 mg, 1500 mg,
1200 mg, 600 mg, 300mg, 150mg
Excipients q.s.

BOLUS

Suspension contains 125mg/ 5ml
Albendazole IP
30ml, 60ml, 90ml,
500ml, 1Ltr, 5 Ltr
Excipients q.s.

LIQUID

Oral Powder contains
5%w/w, 10%w/w, 15%w/w
Albendazole IP
30g, 100g, 300g, 500g, 1 Kg
Excipients q.s.

POWDER

Schedule of Deworming:

- Regular Deworming after an interval of three months in all the species
- Multiple Deworming at the interval of two or three weeks enhances flucidal activity.

Withdrawal Period:

Milk : 7 days; Meat : 14 days

For oral use in animals:

ALBORVET is effective against

- Round worms
- Lung worms
- Tape worms
- Liver flukes

BOLUS

Dosage & Administration

As directed by the veterinarian.

For round worms/ tape worms infestations:

Cattle & Buffalo: 7.5 mg/ kg body weight.

For liver flukes infestation:

Cattle & Buffalo: 15 mg/ kg body weight.

LIQUID

Dosage:

Albendazole

5 - 7.5 mg per kg of body weight

Cattle	: 60 -90 ml
Sheep and Goat	: 2 -3 ml per 10 kg of body weight
Poultry	: 30 -45 ml for 100 Birds
Dogs	: 3 -5 ml twice daily for 3 days

(For liver fluke and tape worms dosage to be increased as per veterinarian's advice)

POWDER

Dosage :

For Roundworms and Tapeworms

Sheep, Goat, Horse, Camel, Pig	- 5mg -3gm per 30kg body weight
Cattle, Buffaloes	- 7.5mg - 15gm per 100kg body weight
Dog, Cat	- 30mg - 12gm per 20kg body weight

For Flukes

Sheep, Goats	- 7.5mg - 9gm per 60kg body weight
Cattle, Buffaloes	- 15mg - 30gm per 100kg body weight

For Poultry

Mixed infection of round worms, Tapeworms and Capillary worms
- 40 to 60gm of Alborvet Powder 5% for 100 birds.