



DOXORVET-N

Doxycycline Hyclate and Neomycin Sulphate Powder

Composition:

Each bolus contains:

Doxycycline IP 160mg
Neomycin IP 400mg

Oral Powder contains:

Doxycycline Hyclate 10% w/w
Neomycin Sulphate IP
Equivalent to Neomycin 10% w/w
Excipients Q.S.

INDICATIONS:

Gastrointestinal respiratory infections caused by Doxycycline and Neomycin sensitive micro-organisms like Bordetella, Campylobacter, Chlamydia, E. Coli, Haemophilus, Mycoplasma, Pasteurella, Ricktsia, Salmonella, Staphylococcus and Streptococcus spp, in calves, goats, poultry, sheep, and swine.

ADMINISTRATION:

Doxorvet N can be administered via clean drinking water in poultry and swine or mixed with feed for calves, goats, and sheep. Mix medicated water and feed properly.

WITHDRAWAL TIMES:

Poultry: meat and egg 7 days.
Cattle, Sheep, Goat, and
Swine: meat 21 days.

DOSAGE:

Poultry & Swine: Prevention and treatment dissolve 1g in 1.5 liters of drinking water for 3-5 consecutive days.
Cattle, Sheep, and Goats: Add 0.1g per 1Kg body weight in feed daily for 3-5 days.
Only medicated water and feed should be consumed during treatment period.
Discard unused medicated water and feed after 24 hours.

CONTRAINDICATIONS:

GRD-Doxyneom should not be administered concurrently with penicillin, cephalosporins, quinolones, cyclserine and other bacteriostatic agents, as over dosage may easily occur. Hypersensitivity reaction may occur in some cases. Administration of GRD-Doxyneom to young and growing animals should be with care and under guidance of a veterinarian.

STORAGE:

Store safely in a cool dark dry place, temperature should not be exceed 25 C.
Always keep the medicine in a closed container when not in use.

WARNINGS:

For animals use only.
Keep out of the reach of children.