



# ENTORVET

## BROAD SPECTRUM ANTIBIOTIC

### Composition

#### Each ml contains

Enrofloxacin IP Vet 100mg  
Aqueous base q.s.

LIQUID

#### Each bolus contains:

Enrofloxacin IP 1500 mg  
Excipients q.s.

BOLUS

### Description

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella and Mycoplasma spp.

### Indications

Gastrointestinal infections, respiratory infections and urinary tract infections caused by enrofloxacin sensitive micro-organisms, like Campylobacter, E.coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp. in calves, goats, poultry, sheep and swine.

### Contra Indications

Hypersensitivity to enrofloxacin  
Administration to animals with a seriously impaired hepatic or renal function.  
Administration to poultry producing eggs for human consumption.  
Administration to pregnant or lactating animals  
Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

### Side Effects

Administration to young animals during growth can cause cartilage lesions in joints. Hypersensitivity reactions.

### Dosage & Administration

#### Poultry

##### For Prevention

2.5mg to 5mg per kg body weight for 3 to 5 days (1 ml of ENTORVET in 4-8 litres of drinking water)

##### For treatment

10mg per kg body weight for 3 to 5 days (1ml of ENTORVET in 2 litres of drinking water) or as directed by the Veterinary Practitioner only

#### Large Animals

2-3ml for 30kg body weight

#### Sheep & Goat

2ml 25kg body weight for 3-5 days or as directed by the veterinarian.

#### Dosage :

Cattle - 2 bolus for 300 kg body weight.

### Storage

Store in a cool and dry place. Protect from direct sunlight and heat.

### Withdrawal Period

Meat : 5-14 days

Milk : 3 days

LIQUID

BOLUS