



FENOVET

A BROAD SPECTRUM DEWORMER

MICRONIZED

Composition:

Each Bolus contains
Fenbendazole IP 3000 mg, 1500mg, 600mg, 150mg
Excipients q.s.

Schedule of Deworming:

- Regular deworming after an interval of three months in all the species

Indication

Infestation with gastrointestinal nematodes, lungworms and tapeworms, such as *Haemonchus* spp., *Ostertagia* spp., *Trichostrongylus* spp., *Cooperia* spp., *Nematodirus* spp., *Bunostomum* spp., *Gaigeria* spp., *Capillaria* spp., *Oesophagostomum* spp., *Chabertia* spp., *Trichuris* spp., *Strongyloides* spp., *Dictyocaulus* spp., *Toxascaris vitulorum*, *Moniezia* spp. Infestation of horses with large and small *Strongylus* spp., *Parascaris equi*, *Oxyuris equi* and *Strongyloides* spp.

Dosage & Administration

BOLUS

Large Animals - 1-2 Bolus /200-400Kg.B.Wt
Sheep,Goat - 1 Tablet/60-120Kg B.Wt.
Dog - 1 Tablet/12Kg B.Wt.(50 mg/Kg.B.Wt)

POWDER

Infestation with gastrointestinal nematodes and lungworms:
5 mg of Fenbendazole per kg body weight
or as advised by consultant Veterinarian.

Withdrawal Period:

Cattle must not be slaughtered within 8 days following treatment.
For dairy cattle, there is no milk withdrawal period

For oral use in animals:

FENOVET is effective against

- Round worms
- Tape worms
- Lung worms

STORE PROTECTED FROM LIGHT